



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

giving therefore the numbers 5, 6, 7, and 8 the relative values thus found, and averaging in the ordinary way, we will find that 6 is the average number of good melons, and therefore $\frac{6}{8}$ or $\frac{3}{4}$ is the probability of selecting a good one.

15. Proposed by F. P. MATZ, M. Sc., Ph. D., Professor of Mathematics and Astronomy, in New Windsor College, New Windsor, Maryland.

Todhunter proposes: "From a point in the circumference of a circular field a projectile is thrown at random with a given velocity, which is such that the diameter of the field is equal to the greatest range of the projectile; prove the chance of its falling within the field, is $C=2^{-1}-2\pi^{-1}(\sqrt{2}-1),=.236+.$ " Is this result perfectly correct as to fact?

Comment on the Solution of Problem 15, by JOHN DOLMAN Jr., Philadelphia, Pennsylvania.

I should not presume to criticise the work of so able and celebrated a mathematician as Professor Matz, did I not consider his solutions of Probability No. 15 vicious in their effects upon the minds of students, striking as they do at the root, not only of the doctrine of mean value and probability, but of the integral calculus itself. "Since the projectiles are *thrown* at random they should *fall* at random," is on a par with, "as an arc varies uniformly the sine varies uniformly," or "because acceleration is constant velocity is constant." The third solution is not clear, but how far any given range the favorable chances can be represented by an area, when the projectiles must fall on the arc of a circle, is difficult to understand. Also in the fifth solution it is stated, "for any range PD' , the projectiles falling on the circular arc DMD' are within the field" and then the angle PAD' is adopted as uniformly varying, without giving any reason for it. However interesting this may be as mathematical legerdemain, its effects are vicious when it is published without proper explanation for in my humble opinion it is more important that your readers learn to reason correctly than that they be taught to integrate ingeniously.

NOTE.—The solution of problem 14 was published without comment for the reason that we considered the solution to be correct, and we confess that we do not yet see the force of Mr. Dolman's argument, though we have not had time to give it much thought.

As to the solutions of problem 15, we hold that the first solution is the only correct solution as that one and that one alone involves the strict literal statement of the problem. It is evident that the number of ways the projectile can be thrown is equal to the surface of a hemisphere whose radius is R . If now we find the surface of that part of this hemisphere any point at which if a projectile be thrown the projectile will fall upon the circular field, diameter R , and then divide this surface by the surface of the hemisphere, the result will be the probability required. This method of solution would have to be accepted by the most critical mind. Professor Matz's first solution involves this principle.—EDITOR.

19. Proposed by F. P. MATZ, M. Sc., Ph. D., Professor of Mathematics and Astronomy in New Windsor College, New Windsor, Maryland.

Find the average area of the circle which is the locus of the middle points of all chords passing through a point taken at random in the surface of a given circle.

I. Solution by G. B. M. ZERR, A. M., Principal of High School, Staunton, Virginia.